

# compass



The Evolution of Fundus Perimetry



---

## Company Profile

CenterVue designs and manufactures highly automated medical devices for the diagnosis and management of ocular pathologies, including those that represent the leading causes of blindness.

Our goal is to design Smartly Simple devices and to provide high-value services that enable Eye Care Specialists to better preserve patients' sight and quality of vision, in particular by detecting preventable diseases; and in so doing dramatically improve their quality of life.

CenterVue is headquartered in Padova, Italy, with the US branch in San Jose, California. CenterVue is present in over 70 countries with its distribution network.

---



---

“It seems evident that the main usage for microperimeters is destined to monitor glaucoma damaged residual functional vision [..] For mainstream ophthalmology many indicators point to the fact that microperimeters may take the lead role played by SAP in the last decades.”

Microperimetry and clinical practice: an evidence-based review;  
Samuel N. Markowitz, MD, FRCSC, Sophia V. Reyes, MD. Canadian Journal  
of Ophthalmology Vol. 48, Issue 5.

---



## The Next Level of Fundus Perimetry

### Compatibility with standard 24-2 visual field testing

As a perimeter, the system offers full compatibility with standard 24-2 visual field testing and contains an age-matched database of retinal sensitivity in normal subjects.



24-2 test performed with Compass


### Superior quality of color and red-free images

As a retinal imager, COMPASS uses a confocal optical design, similarly to SLO systems, to capture color as well as red-free images of superior quality.


In addition, a high resolution live image of the retina obtained using infrared illumination is available throughout the test.



Color image: detail of the ONH



Infrared image




Red-free image: detail of the RNFL

### Retinal Tracking

Retinal tracking is at the heart of Fundus Perimetry.

Infrared images, acquired at the rate of 25 images per second, allow for continuous, automated, tracking of eye movements, with positional accuracy in the 10-20 microns range. Determination of eye movements yields to Fixation Analysis, where the location of the functional site of fixation and its stability are computed. Fixation analysis is unique to Fundus Perimetry. Retinal tracking also yields to active compensation of fixation losses, with perimetric stimuli being automatically re-positioned prior to projection based on the current eye position. This mechanism is critical to reduce test-retest variability and ensure accurate correlation between function (i.e. retinal threshold values) and structure (retinal appearance).

Compensation of eye movements takes place before and during the projection of a certain stimulus. In absence of this mechanism, a normal 2-3 degrees shift in eye position occurring at the time of projection of a certain stimulus would easily produce an artifact in VF results, with a wrong sensitivity being reported at that specific location.




Plot of eye movements during VF testing


### Color confocal imaging

SLO systems are superior to conventional fundus cameras in many ways, as they exploit a confocal imaging principle, which limits the effect of backscattered light from deeper layers and provides enhanced image quality in terms of contrast and resolution. Another major advantage of SLO systems is that they operate with much smaller pupils than non-confocal instruments. At the same time, though, SLO systems do not provide color images, as they typically employ monochromatic laser sources, resulting in black and white or pseudo-color images. Differently from existing SLO systems, Compass uses white light instead of monochromatic lasers, hence providing true color images and offering high fidelity to real retinal appearance. Compass images improve the diagnostic capabilities in the management of glaucoma as they offer:

- no need for pupil dilation
- excellent resolution and contrast
- high quality even in presence of media opacities, such as cataract
- optimized exposure of the ONH



Color image: detail of the ONH




Non-confocal imaging: detail of the ONH

## Fixation Analysis In Glaucoma

Studies have demonstrated abnormal fixation characteristics in patients diagnosed with early Primary Open Angle Glaucoma (POAG) and Advanced Glaucoma without other retinal diseases.

For example, fixation instability has been demonstrated in early and moderate POAG [1], while other studies have reported predominantly eccentric fixation in up to 15% of the studied population with advanced glaucoma [2]. Finally, it is known that fixation stability and its location correlate with visual acuity, in particular the more unstable and eccentric fixation is, the lower visual acuity [3].

Fixation analysis with Compass provides additional, quantitative, parameters for assessing visual function.



- Shi Y, Liu M, Wang X, Zhang C, Huang P (2013) Fixation behavior in primary open angle glaucoma at early and moderate stage assessed by the MicroPerimeter MP-1. Journal of glaucoma 22 (2):169-173.
- Kameda T, Tanabe T, Hangai M, Ojima T, Aikawa H, Yoshimura N (2009) Fixation behavior in advanced stage glaucoma assessed by the MicroPerimeter MP-1. Japanese journal of ophthalmology 53 (6):580-587.
- Morales MU, Saker S, Mehta RL, Rubinstein M, Amoaku WM (2013) Preferred retinal locus profile during prolonged fixation attempts. Can J Ophthalmol 48 (5):368-374.

## Benefits

- High-resolution confocal imaging of the ONH and of the central retina
- Combined structure and function analysis
- Significantly reduced test – retest variability
- Reliable follow-up
- Fully automated operation
- Comprehensive and clear printout
- Operator friendly
- More patient comfort: test can be suspended at any time without data loss

## Features


- Pupillary tracking
- Fixation control using automated high-resolution, 25 Hz, retinal tracking
- Auto-focus
- Fully automated, visual field testing (24-2, 10-2)
- Confocal, 60° imaging
- Color, red-free, infrared imaging
- Patient vocal guidance during VF testing
- Quantitative analysis of Fixation
- Wired/WiFi connectivity
- Touch-screen operated via tablet
- Non-mydratic operation

# How to Read Printouts

COMPASS combines in one page the standard deviation maps typical of SAP with structure-function correlation data as well as fixation analysis.

## Legend

- Patient info
- Examined eye
- Exam info
- Color image of ONH
- Fundus Perimetry (dB) over red-free image
- Deviation maps
- Standard VF map
- Mean Deviation and Pattern Standard Deviation
- Glaucoma Staging System 2
- Bivariate Contour Ellipse Area indices and fixation plot over IR image
- Fixation graph describing amplitude of eye movements vs. time



# compass

The smartly simple approach

COMPASS performs the entire 24-2 test, including capture of infrared, red-free and color retinal images, in a fully automatic way.



The complete process includes:

- automated alignment of the instrument to the patient, using bilateral pupil tracking;
- auto-focus, correcting for the patient's spherical refraction;
- automated acquisition of live, infrared, retinal images;
- automated execution of the VF test;
- automated collection of fixation data throughout the VF test;
- automated compensation of eye movements during VF testing, using retina and pupil tracking;
- automated acquisition of a color retinal image, including auto-exposure;
- the only intervention by the operator during the process is the selection of the center of the ONH.



Ergonomic and motorized chin rest. Improved cleaning ability of the patient rest cushions.

Touch screen interface and high resolution, 2560x1600 pixel display.

The patient push button is designed for improved ergonomics.

Connectors on the back include 3 USB ports and 1 Ethernet port.

Digital joystick is used for manual alignment and focusing.

## Technical specifications\*

### Class and type of applied part

1, B (according to EN 60601-1).

### Fundus Perimetry:

- Projection field: 30° (radius)
- Background luminance: 31.4 asb
- Maximum luminance: 10000 asb
- Dynamic range: 0 - 50 dB
- Stimulus size: Goldmann III (26")
- Stimulus duration: 200 ms
- Threshold tests: 24-2, 30-2, 10-2
- Fixation control: 25 Hz automated retinal tracking
- Foveal threshold testing
- Automatic pupil measurement

### Fundus Imaging:

- Field of view: 60° (diameter)
- Sensor resolution: 5 Mpixel (2592x1944)
- Light source: infrared (825-870 nm) and white LED (440-650 nm)
- Imaging modalities: color, infrared, red-free
- Resolution: 17 microns

### Other features:

- Automatic operation: auto-alignment, auto-focus, auto-retinal tracking, auto-pupil tracking, auto-exposure, auto-capture
- Non-mydratic operation: minimum pupil size 3 mm
- Working distance: 28 mm
- Auto-focus range: -12D to +15D

- Fixation target: programmable, internal
- User interface: Nexus Tablet with 10.1" multi-touch screen
- Connectivity: Wi-Fi and Ethernet
- Printer: any printer compatible with Tablet
- Hard disk: SSD, 240 GB

### Dimensions:

- Weight: 25 Kg
- Size: H 620 X W 590 X D 360 mm

### Electrical requirements:

- Power: 100-240 VAC, 50-60 Hz
- Consumption: 80 W



\* Specifications are subject to change without notice for improvement.

# compass

---

The first fundus perimeter capable of performing standard 24-2 visual field testing and delivering true color confocal images.

---





# compass

Centervue SpA

Via San Marco 9H  
35129 Padova - Italy

Ph: +39 049 7396 147  
Fax +39 049 7396 148

info@centervue.com  
www.centervue.com

Centervue Inc.

92 Bonaventura Drive,  
San Jose CA 95134 - USA

Ph: +1 408 988 8404  
Fax: +1 408 716 3271

infous@centervue.com  
www.centervue.com